

# An Interventional Study on Antibiotic Cement-coated Nails and Cement Beads in the Management of Infected Nonunion of Long Bones

Prabhat Agrawal<sup>1</sup>, SV Sunny Deol<sup>2</sup>, N Rajkumar<sup>3</sup>, V Arul Jothi<sup>4</sup>, SM Murali<sup>5</sup>

## ABSTRACT

**Background:** The present study was aimed to assess the efficacy of antibiotic cement-coated nails and cement beads among the cases of infected nonunion of long bones.

**Materials and methods:** This prospective interventional study was conducted among the cases admitted in the emergency and outpatient department of Orthopedics in Patna Medical College and Hospital, Patna, with infected nonunion of long bones, during the month of June 2010 to August 2012. All cases with an infected nonunion of extremities with implants were included. A total of 44 cases were included. Data analysis was done using SPSS version 17.

**Results:** In this study, out of 44 infected nonunion, 40 were cured of infection, 3 cases had occasional discharge, and 1 case had a continuous discharge. Secondary procedures were done for the 40 cured cases and among them, 37 showed complete achievement of union, and 3 cases had occasional discharge and no case had a continuous discharge.

**Conclusion:** The use of antibiotic cement-coated nails and cement beads among the cases with infected nonunion of long bones significantly reduced the requirement of a further surgical procedure like debridement of the wound site.

**Keywords:** Antibiotic cement-coated nails, Cement beads, Infected long bone.

*TNOA Journal of Orthopedics and Joint Surgery* (2021): 10.5005/jp-journals-10079-1049

## INTRODUCTION

In the past decades, the incidence of infected nonunion shows a steady increase due to a high rate of trauma from road traffic accidents, industrial injuries leading to open fractures, postoperative infection following open reduction and internal fixation (ORIF), and injurious use of antibiotics resulting in resistant strains of bacteria when compared with past when chronic osteomyelitis was the major cause of infected nonunion.<sup>1</sup>

Causes of infected nonunion are usually inherent to the fracture, like an open wound, loss of soft tissue or bone, severe comminution and gross displacement, vascular compromise, deep-seated infection at fracture area, periosteal stripping, etc.<sup>2</sup> In infected nonunion, secondary infections are common and multiple organisms may grow from cultures taken from sinus tracts and open biopsy specimens of surrounding soft tissue and bone.

Traditionally, the treatment strategy has been a two-staged procedure. The first stage was to control infection that occurred at the fracture site and the second stage involved procedures to achieve bone union. The control of infection requires repeated debridement, intravenous antibiotics<sup>3</sup> along external stabilization. Once the infection has been eradicated secondary procedures like the opening of the medullary canal, freshening of fracture ends, proper reduction of the fracture, internal or external stabilization, and cancellous and/or cortical bone grafting are done to achieve union at the fracture site.

Buchholz and Engelbrecht<sup>4</sup> reported that penicillin, erythromycin, and gentamicin incorporated into the cement used to attach total hip joint prostheses, diffused out into surrounding tissues over a period of months, thereby providing prolonged concentrations of local antibiotic. Klemm<sup>5</sup> used gentamicin-impregnated cement into

<sup>1</sup>Department of Orthopaedics, All India Institute of Medical Sciences, Patna, Bihar, India

<sup>2,5</sup>Department of Orthopaedics, Sri Muthukumaran Medical College Hospital, Chennai, Tamil Nadu, India

<sup>3</sup>Dhanalakshmi Srinivasan Medical College and Hospital, Perambalur, Tamil Nadu, India

<sup>4</sup>Department of Ortho, OMS Hospital, Chennai, Tamil Nadu, India

**Corresponding Author:** SM Murali, Department of Orthopaedics, Sri Muthukumaran Medical College Hospital, Chennai, Tamil Nadu, India, Phone: +91 9445373395, e-mail: Muraliorth@gmail.com

**How to cite this article:** Agrawal P, Sunny Deol SV, Rajkumar N, *et al.* An Interventional Study on Antibiotic Cement-coated Nails and Cement Beads in the Management of Infected Nonunion of Long Bones. *TNOA J Orth Joint Surg* 2021;3(2):76–79.

**Source of support:** Nil

**Conflict of interest:** None

beads and used these to temporarily fill in the dead space created after the debridement of infected bone.

Single-staged procedures such as debridement and application of Ilizarov fixator<sup>6,7</sup> or use of antibiotic cement-impregnated intramedullary nails (ACIINs)<sup>8,9</sup> have been described. The Ilizarov ring fixator has been used after debridement for bone transport or corticotomy distraction after acute docking. This procedure is technically demanding and has significant complications and is best suited for large segmental bone defects<sup>6,7,10,11</sup>

Antibiotic cement-coated nail and antibiotic cement beads are considered as one of the options for treating the infected long bones and also provide better stability to the injured long bone.<sup>8,9</sup>

Use of antibiotic cement-coated nail can allow early weight-bearing and also it reduced the complications associated with fixator, such as infection of the pin tract, stiffness of the joints, etc. Hence, this study was conducted to assess the efficacy of antibiotic cement-coated nails and cement beads in cases of infected nonunion of long bones.

## MATERIALS AND METHODS

This prospective interventional study was conducted among the cases admitted in the emergency and outpatient department of Orthopedics in Patna Medical College and Hospital, Patna with infected nonunion of long bones, during the month of June 2010 to August 2012. All cases with an infected nonunion of extremities with implants in situ were also included in the study. Patients with non-infected nonunion, open physis, infected nonunion with bone gap >4 cm after debridement, and patients with diabetes mellitus, tuberculosis, and HIV were excluded from the study. Hence, a total of 44 cases were included in this study.

The study was approved by the ethical committee of this institution. Informed consent was obtained from the patients, before starting the study. A thorough history was taken to determine if the patient has a history of allergy to antibiotics so that those antibiotics were not used when mixing it with the bone cement. The evaluation of intraoperative culture and sensitivity results guides the selection of the appropriate postoperative intravenous antibiotics.

Preoperative radiographs were taken which helps the operating surgeon to assess the size of the intramedullary nail needed.

The length of the intramedullary nail is determined as follows:

- For K-nail—from the tip of the greater trochanter to the upper border of patella plus 2 cm.
- For V-nail from the tip of the upper end of the fibula to medial malleolus minus 2 cm.
- The diameter of both K-nail and V-nail is selected 2 mm less than the maximum size reamer used as a further 2 mm is makeup by the coating of cement.

All patients were administered 3 weeks of intravenous antibiotics and 3 weeks of oral antibiotics based on their culture and sensitivity reports.

### Procedure

After selecting the patients based upon the aforesaid criteria, the patients were taken for operation after proper counseling and pre-anesthetic check-up if needed. Implant removal was performed first. The sinus tracts were injected with methylene blue and were excised till the bone. The fracture site was opened and radical debridement was performed with excision of the infected bone end, scarred soft tissue, and granulation tissue. The intramedullary canal was reamed to the size 2 mm more than the previous nail size or till the fresh bleeding bone was reached. The reaming and granulation tissue was sent for culture and sensitivity tests. The wound and entire medullary canal were cleaned with 4–5 L of normal saline. Once debridement was completed, the instruments that were used for the dirty portion of the procedure were removed and the patient's limb was prepared again and draped. The surgeon and the rest of the operating team change all gowns and gloves. The clean portion of the procedure then begins, usually by the surgeon inserting the antibiotic cement-coated nail.

Antibiotic impregnated cement-coated nail was prepared, K-nail for the femur, V-nail for tibia, K-wire for the humerus, and radius was used. We used 2 g of vancomycin and 2 g of gentamicin for every 40 g of bone cement. A higher mix ratio will alter the settling property as also the handling property. An intramedullary nail size 2 mm less than the largest reamer size used, with a 7- or 8-mm nail was used most commonly. Manual mixing of cement was performed and cement was applied to the nail in a uniformed fashion. The eye of the nail was kept open for easy removal of the nail. Uniform width was achieved by repeatedly passing it from the nail width measuring scale. A fairly smooth surface is obtained by manually rolling and repeatedly checking of width by passing through nail width measuring scale. The nail was kept for 15 minutes in the air for evaporation of monomer.

For beads—the mixture of antibiotic and PMMA and methylmethacrylate is molded or rolled to 3–10 mm spheres and then strung on to S–S wire no. 20. Antibiotic impregnated cement-coated nails and beads were used for infected nonunion. Retrograde insertion of the nail was done through fracture site for the femur and antegrade insertion for the tibia. In cases with bone defects, approximation of bone ends was attempted. In cases where the fracture site was opened antibiotic-impregnated cement beads were left. Wound cleaning was done and closed in a layer without drain.

### Postoperative Protocol

Removal of stitches was done on the 14th postoperative day. Gentle joint mobilization exercises and muscle strengthening were begun as soon as the cases were comfortable and as tolerated. Gradual weight-bearing generally was given after >2 cortex union achieved. All the cases were followed up at 2, 4, 12, 24 weeks, and then 4 weekly for a minimum period of 2 years. Cement beads and nails were removed after 6–12 weeks. Union is assessed by using Stans et al. scale.<sup>12</sup> The patient were evaluated clinically and radiographically using Stans et al. scale for grading callus formation as a useful indicator of fracture healing (Table 1).

Data analysis was done using a statistical package for social sciences (SPSS)–17.

## RESULTS

In the present study, the majority of the study participants were in the age group of 36–45 years and the proportion of male participants was comparatively slightly more than female participants. The open fracture was the commonly reported fracture followed by closed fracture and pathological fractures. Also, the femur was the commonly affected bone followed by the tibia, humerus, and radius. Methicillin-resistant *Staphylococcus aureus* (MRSA) was the common organism isolated followed by *Pseudomonas*, *Escherichia coli*, and *Klebsiella*. In the present study,

**Table 1:** Grading of callus formation

Grade 0	No identifiable fracture healing
Grade I	Primary bone healing with little or no periosteal new bone formation
Grade II	Periosteal new bone formation on two sides of the bone
Grade III	Periosteal new bone formation on three or four sides of the bone

90.9% of cases had complete eradication of infection, 6.8% of cases had occasional discharge, and 2.3% of cases had continuous wound discharge (Table 2). Secondary procedures were done for the 40 cured cases and among them, 37 showed complete achievement of union, and 3 cases had occasional discharge and no case had a continuous discharge. Thus, the outcome after the second procedure was 92.5% of cases had complete healing and 7.5% of cases had occasional wound discharge (Table 3).

## DISCUSSION

Complete debridement, firm fixation, and the use of high-dose antibiotics are the recommended treatment for infected nonunion of the long bone.<sup>1,6,13,14</sup> Several procedures have been tried for the

better management of the same and intramedullary devices were used in both primary and secondary stages of infection control<sup>8,9</sup> and bone healing,<sup>15</sup> respectively, with remarkable outcomes.<sup>16</sup>

Sinus tract culture reports were negative in some of the studies, where they reported it to have poor diagnostic efficacy.<sup>17-19</sup> Intramedullary infection is the cause for the nonunion of the fracture is the common complication of ORIF either by intramedullary nailing.<sup>20</sup> Open fractures have a higher incidence than closed fractures treated with intramedullary nailing.<sup>20,21</sup> Our series had 20 cases of infected nonunion of which 56.8% of cases had an open fracture, 27.3% of cases had closed fractures, and 15.9% of cases had a pathological fracture due to chronic osteomyelitis at the time of initial injury. At the time of presentation, all the cases of nonunion were due to postoperative infection except one which was pathological fracture due to chronic osteomyelitis. May et al.<sup>22</sup> classified bone defect into <6 and >6 cm, while Jain and Sinha<sup>13</sup> proposed a classification of the defect into <4 and >4 cm. We do not include the cases having bone defects >4 cm, which may be the reason for the good outcome of our study.

In this study, in most of the cases organism isolated from culture and sensitivity report of granulation/necrotic tissue obtained during debridement was MRSA, which was found to be reported in 59.1% of cases followed by *Pseudomonas* in 22.7%, *E. coli* in 15.9%, and *Klebsiella* in 2.3% of cases. These findings were compared with the findings of Shyam et al.<sup>23</sup> who reported that out of 25 cases, *Staphylococcus aureus* was found in 22 cases. The available studies thus show that MRSA is the most common causative organism of infected nonunion. Thonse and Conway<sup>24</sup> reported that MRSA was found in 57.69% of cases followed by other bacteria. The important part of this study is to analyze the prime role of the antibiotic cement-coated nails and cement beads in the control of infection. In this study, a total of 44 cases were treated with antibiotic cement-coated nails and cement beads, at first thorough debridement done and antibiotic cement-coated nail inserted intramedullary and antibiotic cement beads under

**Table 2:** Characteristics and outcome after the first procedure

	Frequency	Percentage
Age groups		
16–25 years	09	20.5
26–35 years	13	29.5
36–45 years	22	50
Sex		
Male	24	54.5
Female	20	45.5
Side affected		
Right extremity	25	56.8
Left extremity	19	43.2
Type of fracture		
Open fracture	25	56.8
Closed fracture	12	27.3
Pathological fracture	07	15.9
Bone affected		
Femur	23	52.3
Tibia	16	36.4
Humerus	03	6.8
Radius	02	4.5
Organism isolated		
Methicillin-resistant <i>Staphylococcus aureus</i>	26	59.1
<i>Pseudomonas</i>	10	22.7
<i>Escherichia coli</i>	07	15.9
<i>Klebsiella</i>	01	2.3
Outcome		
Complete eradication of infection	40	90.9
Occasional discharge	03	6.8
Continuous discharge	01	2.3
Time taken for complete healing (weeks) (N = 40)		
4–6 weeks	20	50
6–8 weeks	15	37.5
8–10 weeks	03	7.5
10–12 weeks	02	05
Grading of callus formation		
Grade 0	04	9.1
Grade I	15	34.1
Grade II	19	43.2
Grade III	06	13.6

**Table 3:** Outcome after the second procedure

Secondary procedure	Frequency	Percentage
Exchange nailing	18	45
Illizarov fixator	09	22.5
Orthofix	06	40
Plating with bone graft	07	17.5
Outcome after the secondary procedure		
Achieving union	37	92.5
Occasional discharge	03	7.5
Time taken for complete healing (weeks) (N = 37)		
28–38 weeks	10	27.0
38–48 weeks	11	29.7
48–58 weeks	06	16.2
58–68 weeks	05	13.5
68–78 weeks	03	8.1
78–88 weeks	01	2.7
88–98 weeks	01	2.7
Complications		
Joint stiffness	10	25
Cement nail rebonding	06	15
Shortening	06	15

subcutaneous tissue overlying fracture site, and skin was closed primarily without drain in 70% of cases (the cases includes open fracture and pathological fracture) and in remaining 30% of cases only antibiotic cement-coated nail was used and an incision was closed primarily without drain.

In the present study, we found that out of 44 cases, 40 cases (90.9%) were completely cured of infection, 3 cases (6.8%) had occasional discharge, and 1 case (2.3%) had a continuous discharge. The mean duration of infection control was 7.4 weeks. Those and Conway<sup>24</sup> had studied antibiotic cement-coated interlocking nails for the management of infected nonunions and segmental bone defects in cases of infected nonunion with bone defects in 20 patients. They reported infection control in 95% of their study participants. A study conducted by Shyam et al.<sup>23</sup> reported that all their study cases achieved infection control by using antibiotic-coated nails. From these studies, an inference can be made that antibiotic cement-coated nails and cement beads are very effective in controlling infection at the fracture site in cases of infected nonunion with bone defect <4 cm.

In the present study, we achieved infection control in 40 patients (90.9%). Among them, additional procedure in the form of exchange nailing with the interlocking nail was done in 18 cases, Ilizarov fixator in nine cases, orthofix in six cases, and plating with bone graft was done in seven cases. After which union was achieved in 37 cases (92.5%) out of 40 cases. The mean time for the radiological union was 43.6 weeks. The remaining three (7.5%) cases in which we did not achieve union were required additional debridement, curettage, and refixation with bone grafting but the result of these additional procedures was not possible to add in this study because of the short duration of the study. This study shows that antibiotic cement-coated nails were very effective in controlling infection in cases of infected nonunion but could not provide absolute stability to fracture so additional surgery was done to achieve bony union.

The limitation of our study is that we had selected cases with bone defects <4 cm only and we had not used cement-coated interlocking nails in our study.

## CONCLUSION

At the end of our study, we conclude that infected nonunion of long bones can be best treated with the use of antibiotic-coated intramedullary nails and cement beads. Also, it reduces the requirement of wound debridement and provides better stability to the fracture. Duration of control of infection in such cases can be drastically reduced with help of these methods. In some cases without significant bone defects, we can achieve bony union without the need for secondary procedures. Secondary procedures are generally needed to achieve osseous union.

## REFERENCES

- Lee SJ, Jang DW, Kim TS, et al. Treatment of chronic osteomyelitis by saucerization and gentamicin-PMMA beads. *J Korean Orthop Assoc* 1989;24(2):549–556. DOI: 10.4055/jkoa.1989.24.2.549.
- Court-Brown CM. Fractures of the tibia and fibula. In: Buchholz RW, Heckman JD, Court-Brown CM, ed. *Rockwood and Green's fractures in adults*. 6th ed., Lippincott Williams and Wilkins; 2006. pp. 2080–2146.
- Patzakis MJ, Zalavras CG. Chronic posttraumatic osteomyelitis and infected nonunion of the tibia: current management concepts. *J Am Acad Orthop Surg* 2005;13(6):417–427. DOI: 10.5435/00124635-200510000-00006.
- Buchholz HW, Engelbrecht H. Über die depotwirkung einiger antibiotica bei vermischung mit dem kunstharz palacos. *Chirurg* 1970;41(11):511–515.
- Klemm K. Gentamycin-PMMA-kugeln in der behandlung abszedierender knochen- und weichteilinfektionen. *Zentralbl Chir* 1979;104(14):934–942.
- Toh CL, Jupiter JB. The infected nonunion of the tibia. *Clin Orthop Relat Res* 1995;315(315):176–191. DOI: 10.1097/00003086-199506000-00019.
- Wu CC, Shih CH. Distal tibial nonunion treated by intramedullary reaming with external immobilization. *J Orthop Trauma* 1996;10(1):45–49. DOI: 10.1097/00005131-199601000-00007.
- Paley D, Herzenberg JE. Intramedullary infections treated with antibiotic cement rods: preliminary results in nine cases. *J Orthop Trauma* 2002;16(10):723–915. DOI: 10.1097/00005131-200211000-00007.
- Qiang Z, Jun PZ, Jie XJ, et al. Use of antibiotic cement rod to treat intramedullary infection after nailing: preliminary study in 19 patients. *Arch Orthop Trauma Surg* 2007;127(10):945–951. DOI: 10.1007/s00402-007-0315-x.
- Maini L, Chadha M, Vishwanath J, et al. The Ilizarov method in infected nonunion of fractures. *Injury* 2000;31(7):509–517. DOI: 10.1016/s0020-1383(00)00036-x.
- Song HR, Cho SH, Koo KH, et al. Tibial bone defects treated by internal bone transport using the Ilizarov method. *Int Orthop* 1998;22(5):293–297. DOI: 10.1007/s002640050263.
- Stans AA, Morrissy RT, Renwick SE. Femoral shaft fracture treatment in patients age 6 to 16 years. *J Pediatr Orthop* 1999;19(2):222–228. DOI: 10.1097/01241398-199903000-00017.
- Jain AK, Sinha S. Infected nonunion of the long bones. *Clin Orthop Relat Res* 2005;431:57–65. DOI: 10.1097/01.blo.0000152868.29134.92.
- Struijs PA, Poolman RW, Bhandari M. Infected nonunion of the long bones. *J Orthop Trauma* 2007;21(7):507–511. DOI: 10.1097/BOT.0b013e31812e5578.
- Shahcheraghi GH, Bayatpoor A. Infected tibial nonunion. *Can J Surg* 1994;37(3):209–213.
- Eshima I, Feibel RJ, Louie KW, et al. Combined muscle flap and Ilizarov reconstruction for bone and soft tissue defects. *Clin Orthop Relat Res* 1996;332(332):37–51. DOI: 10.1097/00003086-199611000-00007.
- Donati D, Biscaglia R. The use of antibiotic-impregnated cement in infected reconstructions after resection for bone tumours. *J Bone Joint Surg Br* 1998;80(6):1045–1050. DOI: 10.1302/0301-620x.80b6.8570.
- Zuluaga AF, Galvis W, Jaimes F, et al. Lack of microbiological concordance between bone and non-bone specimens in chronic osteomyelitis: an observational study. *BMC Infect Dis* 2002;16(1):2–8. DOI: 10.1186/1471-2334-2-8.
- Akinyoola AL, Adegbehingbe OO, Aboderin AO. Therapeutic decision in chronic osteomyelitis: sinus track culture versus intraoperative bone culture. *Arch Orthop Trauma Surg* 2009;129(4):449–453. DOI: 10.1007/s00402-008-0621-y.
- Joshi D, Ahmed A, Krishna L, et al. Unreamed interlocking nailing in open fractures of tibia. *J Orthop Surg (Hong Kong)* 2004;12(2):216–221. DOI: 10.1177/230949900401200215.
- Coles CP, Gross M. Closed tibial shaft fractures: management and treatment complications: a review of the prospective literature. *Can J Surg* 2000;43(4):256–262.
- May JW, Jupiter JB, Weiland AJ, et al. Clinical classification of post-traumatic tibial osteomyelitis. *J Bone Joint Surg Am* 1989;71(9):1422–1428. DOI: 10.2106/00004623-198971090-00027.
- Shyam AK, Sancheti PK, Patel SK, et al. Use of antibiotic cement-impregnated intramedullary nail in treatment of infected non-union of long bones. *Indian J Orthop* 2009;43(4):396. DOI: 10.4103/0019-5413.55468.
- Thonse R, Conway J. Antibiotic cement-coated interlocking nail for the treatment of infected nonunions and segmental bone defects. *J Orthop Trauma* 2008;21(4):258–688. DOI: 10.1097/BOT.0b013e31803ea9e6.